




swipe to
see more



PRE-ECLAMPSIA AWARENESS MONTH




AS TOLD BY OUR
CONSULTANT
OBSTETRICIAN
MR. DANIEL STOTT



LONDON
PREGNANCY
CLINIC





swipe to
see more



WHAT?

WHAT IS PRE-ECLAMPSIA



HYPERTENSIVE DISORDER


PERSISTENT HIGH BLOOD
PRESSURE THAT DEVELOPS DURING
PREGNANCY (**USUALLY FROM 20
WEEKS**) OR THE POSTPARTUM
PERIOD (**UP TO 6 WEEKS
POSTPARTUM**)

COMPLICATIONS

WITH CAREFUL MANAGEMENT
MOST PREGNANCIES WITH
PRE-ECLAMPSIA WILL HAVE
GOOD OUTCOMES, BUT
COMPLICATIONS INCLUDE
PROBLEMS WITH THE GROWTH
OF THE BABY, KIDNEY & LIVER
DAMAGE.



swipe to
see more



KEY STATS?

- **HYPERTENSIVE DISORDERS OF PREGNANCY (HDP) AFFECT 5-10% ALL PREGNANCIES WORLDWIDE**
- **APPROXIMATELY 76,000 WOMEN & 500,000 BABIES DIE EACH YEAR WORLDWIDE DUE TO PRE-ECLAMPSIA**



**OF MATERNAL DEATHS DUE
TO PRE-ECLAMPSIA ARE
PREVENTABLE**



swipe to
see more



SYMPTOMS?

PROBLEMS
WITH EYE SIGHT

HEADACHES


NAUSEA
& VOMITING

SHORTNESS
OF BREATH

SUDDEN
WEIGHT GAIN


STOMACH OR
ABDOMEN PAIN

SWELLING
IN HANDS &
FACE





swipe to
see more



MANAGEMENT

HDP CAN DEVELOP FROM 20 WEEKS OF PREGNANCY OR UP TO 6 WEEKS AFTER CHILDBIRTH. HIGH BLOOD PRESSURE CAN USUALLY BE CONTROLLED BY ANTIHYPERTENSIVE MEDICATION, & THE RISK OF PRE-ECLAMPSIA IN PREGNANCY CAN BE REDUCED BY MAINTAINING A HEALTHY LIFESTYLE.*



*PLEASE NOTE: ALL OF THESE MAY HELP DECREASE RATES OF PRE-ECLAMPSIA, BUT SHOULD BE CONSULTED WITH A HEALTHCARE PROFESSIONAL PRIOR TO ACTION.

thank you
for reading

• WHAT WE OFFER

MR. DAN STOTT LEADS LONDON
PREGNANCY CLINIC'S OBSTETRIC
CONSULTATIONS

➤ HE HELPS PROVIDE
MOTHERS WITH
POST-NATAL CARE

➤ HE CAN ADDRESS ANY
CONCERNS MOTHERS
HAVE DURING OR
POST PREGNANCY

QUALIFICATIONS:
BA(HONS) 1ST CLASS, KCL
MBBS, ST GEORGE'S, UCL
MRCOG, 2017
MD (RESEARCH THESIS), KCL

